

For more information, contact:

Lauren P. Riley

847-384-4031

708-227-1773

pearson@aaos.org

Kayee Ip

847-384-4035

312-543-3211

ip@aaos.org

Vitamin D deficiency and insufficiency impacts children's risk for severe forearm fractures

NEW ORLEANS, La. (March 6, 2018)—Children who are vitamin D deficient have a greater risk of having more severe forearm fractures requiring surgical treatment, according to a new study presented today at the 2018 Annual Meeting of the [American Academy of Orthopaedic Surgeons](#) (AAOS). This is the first report that shows the important link between low vitamin D levels and the severity of fractures in children caused by low-energy, less traumatic events such as falling off a bike or falling while running.

Fractures in children are very common, with some estimates as high as 50 percent of boys and 40 percent of girls having at least one fracture by age 18. Of these fractures, the forearm is the most common site, accounting for approximately 25 percent of all pediatric fractures in the U.S.

“Not only are forearm fractures common in children, but so is vitamin D deficiency and insufficiency,” said Pooya Hosseinzadeh, MD, assistant professor, Department of Orthopaedic Surgery at Washington University School of Medicine in St. Louis. “Knowing that vitamin D deficiency can lead to negative calcium balance, low bone mineral density and quality leading to compromised bone strength, it makes sense for patients to be more susceptible to fractures at lower impact load and more susceptible to greater severity when fractures do occur.”

In this [study](#), 100 children (ages 3 to 15; 65 percent male, 35 percent female) with low-energy forearm fractures were prospectively enrolled. Each participant filled out a questionnaire focusing on risk factors for vitamin D deficiency. The mean 25-hydroxyvitamin D (25[OH]D) concentration was 27.5 ± 8.3 ng/ml.

The fractures were then categorized as requiring non-operative or operative management. The children's vitamin D status was based on measurement of 25(OH)D concentration obtained during the clinic visit, and was compared between the two fracture groups.

Among the results of the study:

- Using [Endocrine Society guidelines](#), 21 percent of patients were vitamin D deficient (25(OH)D \leq 20 ng/ml) and 49 percent had vitamin D insufficiency (25(OH)D: 21 - 29 ng/ml).
- There was a dramatic difference in incidence rates of vitamin D deficiency between those treated surgically (50 percent) compared to those in the non-operative group (17 percent).
- Being vitamin D deficient was also associated with a greater risk of needing operative management (28.6 percent versus 7.6 percent non-operative).
- Being overweight/obese and nonwhite increased the likelihood of vitamin D deficiency.
- Seventy-five percent of children in the operative group were obese or overweight, compared to only 32 percent of children in the non-operative group.
- Patients requiring operative management were older than 10 years of age and had greater BMI than patients not requiring surgery.

“This study provides an important takeaway for parents and pediatricians,” explains Dr. Hosseinzadeh. “If a child does have a forearm fracture, we would encourage the physician to check the patient's vitamin D levels. The good news is that in most cases, children can reduce deficiency with a vitamin D supplement and increasing outdoor activity.”

[2018 AAOS Annual Meeting Disclosure Statement](#)

The American Academy of Orthopaedic Surgeons

With more than 38,000 members, the [American Academy of Orthopaedic Surgeons](#) (AAOS) is the world's largest medical association of musculoskeletal specialists. The AAOS provides education programs for orthopaedic surgeons and allied health professionals, champions and advances the highest quality musculoskeletal care for patients, and is the authoritative source of information on bone and joint conditions, treatments and related issues.

Visit AAOS at:

[Newsroom.aaos.org](#) for bone and joint health news, stats, facts, images and interview requests.

[ANationinMotion.org](#) for inspirational patient stories, and orthopaedic surgeon tips on maintaining bone and joint health, avoiding injuries, treating musculoskeletal conditions and navigating recovery.

[Orthoinfo.org](#) for patient information on hundreds of orthopaedic diseases and conditions.

[Facebook.com/AAOS1](#)

[Twitter.com/AAOS1](#)

[Instagram.com/AAOS_1](#)